

Analysis of Algorithms — Tutorial

Problem 5-1

Given an array a of length n , an algorithm compares all pairs $(a[i], a[j])$ for all $i < j \leq n$, and then calls itself recursively on all proper prefixes of a .

How often does the algorithm compare two pairs? Use the repertoire method!

Solution: The recurrence is $R_0 = 0$ and

$$R_n = \binom{n}{2} + \sum_{k=0}^{n-1} R_k$$

for $n \geq 1$. Testing a couple of values for R_n , we obtain the repertoire:

R_0	R_n	$f_n = R_n - \sum_{k=0}^{n-1} R_k$
1	a^n	$\frac{2-a}{1-a}a^n - \frac{1}{1-a}$
1	2^n	1
1	1	$1-n$
0	n	$n - \binom{n}{2}$

To get $\binom{n}{2}$, we can use the last line. To get rid of the linear and constant factors, we use the third and finally the second line, and obtain

$$-\left(n - \binom{n}{2}\right) - (1-n) + 1 = \binom{n}{2}.$$

Fortunately, this also holds for $R_0 = 0$, and the solution is

$$R_n = 2^n - n - 1.$$

Problem 5-2

Compute the number of iterations of the **while**-loop for $0 < i$ and arbitrary j .

```

while  $i \leq j$ 
     $i := i + j$ ;
    if  $i > j$  then  $j := j + 10$ ;
    
```

Solution:

Let i_n and j_n be the numbers i, j after the n th iteration of the while-loop. For $i_0 > j_0$ there are no iterations. Let therefore $0 < i_0 \leq j_0$ in the following. We get the recurrence

$$\begin{aligned}i_n &= i_{n-1} + j_{n-1} \\j_n &= j_{n-1} + 10.\end{aligned}$$

This immediately yields

$$\begin{aligned}j_n &= j_0 + 10n \\i_n &= i_{n-1} + 10(n-1) + j_0 \\&= i_0 + \sum_{k=1}^n (10(k-1) + j_0) \\&= i_0 + 5n(n-1) + nj_0.\end{aligned}$$

The **while**-loop is iterated while $i_n - j_n \leq 0$ holds, i.e., as long as

$$5n^2 + (j_0 - 15)n + i_0 - j_0 \leq 0.$$

For positive n this inequality holds while

$$n \leq \frac{15 - j_0 + \sqrt{(j_0 - 15)^2 - 20(i_0 - j_0)}}{10} =: a(i_0, j_0).$$

Therefore, the **while**-loop is iterated $\lfloor a(i_0, j_0) \rfloor + 1$ times.

Homework Assignment 5-1 (10 Points)

Use summation factors to solve the following recurrence:

$$\begin{aligned}a_0 &= 0 \\a_n &= \frac{a_{n-1}}{n} + \frac{1}{(n-1)!} \quad \text{for } n \geq 1\end{aligned}$$

Solution: Plugging $y_n = 1/(n-1)!$ and $x_n = 1/n$ into the formula known from the lecture yields:

$$\begin{aligned}a_n &= \frac{1}{(n-1)!} + \sum_{j=1}^{n-1} \frac{1}{(j-1)!} \frac{1}{j+1} \cdots \frac{1}{n} \\&= \frac{1}{(n-1)!} + \sum_{j=1}^{n-1} \frac{j}{n!} \\&= \frac{1}{(n-1)!} + \frac{1}{n!} \frac{n(n-1)}{2} \\&= \frac{2n}{2n!} + \frac{n(n-1)}{2n!} \\&= \frac{n+1}{2(n-1)!}\end{aligned}$$

Homework Assignment 5-2 (10 Points)

Use the repertoire method to find a closed form for the following recurrence:

$$\begin{aligned}a_0 &= 5 \\a_1 &= 9 \\a_n &= na_{n-1} + n^2a_{n-2} - n^4 - 3n^2 + 5 \quad \text{for } n \geq 2\end{aligned}$$

Solution: Let $f(n) = -n^4 - 3n^2 + 5$, i.e., $f(n) = a_n - na_{n-1} - n^2a_{n-2}$.

a_n	$f(n)$	a_0	a_1
1	$-n^2 - n + 1$	1	1
n	$-n^3 + n^2 + 2n$	0	1
n^2	$-n^4 + 3n^3 - n^2 - n$	0	1

Let Z_i for $i = 1, 2, 3$ be the solutions of the first, second, and third line, respectively. Then $f(n) = 5Z_1 + 3Z_2 + Z_3$. For these, a_0 and a_1 are correct, and thus $a_n = 5 \cdot 1 + 3n + n^2$.