**RWTH Aachen University** **WS 2010/11**
**Lehrgebiet Theoretische Informatik** Tutorial 12
Kneis—Langer—Sikdar 18.01.2011

# Tutorial Exact Algorithms

**Exercise T25** An $n \times n$ matrix is *stochastic* if all its entries are nonnegative and every row sums up to 1. An $n \times n$ matrix is *doubly stochastic* if its entries are nonnegative and each row and each column sums up to 1.

1. Let $M$ be the incidence matrix of a regular bipartite graph of degree $r \geq 1$. Show that $\frac{1}{r}M$ is doubly stochastic.
2. The Permanent Inequality states that for any doubly stochastic matrix $A$

$$\operatorname{perm}(A) \geq \operatorname{perm}\left(\frac{1}{n}J_n\right) \geq \frac{n!}{n^n},$$

where $J_n$ is the all-ones $n \times n$ matrix. This inequality was conjectured by van der Waerden in 1926 and proved independently by Egorichev and Falikman in 1980. Use this inequality to show that a regular bipartite graph of degree $r$ has at least $(r/e)^n$ perfect matchings.

**Exercise T26** The Hamiltonian Cycle Problem. Given a graph $G = (V, E)$ decide whether $G$ has a simple cycle $(u_1, u_2, \ldots, u_{|V|}, u_1)$ containing all vertices. Formulate this problem as an Inclusion-Exclusion formula and show that it can be solved in time $O^*(2^n)$ and polynomial space.

**Solution** Fix a vertex $s \in V(G)$ and assume that $V = \{s, u_1, \ldots, u_n\}$. In order to formulate this problem as an IE-formula, we need to decide what the objects are and what properties they must satisfy. An object in this context is a closed walk that starts and ends at $s$ and is of length $n+1$. An object satisfies property $A_v$ for $v \in \{u_1, \ldots, u_n\}$ if it contains vertex $v$. Clearly an object that satisfies all properties $A_{u_1}, \ldots, A_{u_n}$ is a Hamiltonian cycle of $G$. Now

$$|A_{u_1} \cap \cdots \cap A_{u_n}| = \sum_{X \subseteq V \setminus s} (-1)^{|X|} \left| \bigcap_{i \in X} \bar{A}_{u_i} \right|.$$

The term $\left|\bigcap_{i \in X} \bar{A}_{u_i}\right|$ represents the number of closed walks of length $n+1$ that start and end at $s$ and avoid the vertices in $X$. This can be easily computed by finding the $(s,s)$th entry in the $(n+1)$st power of the adjacency matrix of $G \setminus X$. The time taken to compute this for a fixed subset $X$ is $O(n^{\omega+1})$ and the space required is $O(n^2)$. An alternative way of computing the terms $\left|\bigcap_{i \in X} \bar{A}_{u_i}\right|$ is by dynamic programming. For $X \subseteq \{u_1, \ldots, u_n\}$ and $u \in V \setminus X$ define $P_X(u, k)$ to be the number of walks from $s$ to $u$ of length $k$ that avoid all the vertices in $X$. Then

$$\left| \bigcap_{i \in X} \bar{A}_{u_i} \right| = \sum_{u \in N(s) \setminus X} P_X(u, n).$$

Also $P_X(u,0) = 1$ if $u = s$ and 0 otherwise. For $k \geq 1$,

$$P_X(u,k) = \sum_{v \in N(u) \setminus X} P_X(v, k-1).$$

For a fixed $X \subseteq \{u_1, \ldots, u_n\}$, the time taken to compute $\left|\bigcap_{i \in X} \bar{A}_{u_i}\right|$ is $O(mn)$ and the space required is $O(n^2)$. Thus the total time taken to evaluate $|A_{u_1} \cap \cdots \cap A_{u_n}|$ is $O^*(2^n)$ and the space required is polynomial.

**Homework Assignment H26 (10 Points)** Given an $n \times n$ matrix $A = [a_{ij}]$ show that

$$\operatorname{perm}(A) = \sum_{S \subseteq [n]} (-1)^{|S|} \prod_{i=1}^{n} \sum_{j \notin S} A[i,j].$$

**Hint.** One can view the elements of $A$ as variables and the permanent of $A$ as a multinomial polynomial. One way of proving the above is by determining the coefficient of these terms on either side of the purported identity. For example, for a fixed permutation $\pi\colon [n] \to [n]$, the coefficient of the term $a_{1,\pi(1)}, \ldots, a_{n,\pi(n)}$ is a 1 on either side. Now determine the coefficient of terms that involve variables from $k$ columns. The coefficient of such a term is 0 on the left-hand side. What is the coefficient on the right-hand side? Conclude that the permanent of an $n \times n$ matrix can be computed in time $O^*(2^n)$ and space polynomial in $n$.

**Homework Assignment H27 (10 Points)** The Travelling Salesperson Problem (TSP). Given a graph $G = (V, E)$ where $V = \{s, u_1, \ldots, u_n\}$ and a weight function $w\colon E \to \{1, \ldots, c\}$, determine the weight of the smallest tour that starts and ends at $s$. Use Inclusion-Exclusion to solve TSP in time $O^*(2^n c)$. What is the space requirement of your algorithm? Comment on whether it is a polynomial space algorithm.