RWTH Aachen University **WS 2010/11**
Lehrgebiet Theoretische Informatik Tutorial 14
Kneis—Langer—Sikdar 01.02.2011

Tutorial Exact Algorithms

The hints for the exercises and assignments are to found in the slides for the lectures.

Exercise T29 Properties and uses of the Convolution Product. Let U be a set of n elements and let f, g, h be functions from 2^U to $\mathbf{Z}$. Recall that the convolution of f and g, denoted by $f * g$, is a function from 2^U to $\mathbf{Z}$ defined as follows. For all $S \subseteq U$,

$$(f * g)(S) = \sum_{\substack{X,Y \subseteq S \\ X \cap Y = \emptyset}} f(X)g(Y).$$

Show that the convolution product is commutative and associative, that is,

$$\begin{aligned} f * g &= g * f. \\ (f * g) * h &= f * (g * h). \end{aligned}$$

Given a graph $G = (V, E)$, define $f \colon 2^V \to \{0, 1\}$ as follows. For $S \subseteq V$, we let $f(S) = 1$ iff S is an independent set in G.

1. Show that $(f * f)(V)$ represents the number of two colorings of the graph G.
2. What does the iterated k-fold convolution $(f * \cdots * f)(V)$ (k times) represent? How would you compute the chromatic number of a graph G, if you were given an algorithm that computes the convolution $f * g$?
3. If $f(S)$ is defined to be 1 iff S is a dominating set of G, then what does the iterated k-fold convolution $(f * \cdots * f)(V)$ (k times) represent?

Exercise T30 Fast Ranked Zeta Transform. In class we saw how to compute the zeta and Möbius transforms of a function $f \colon 2^U \to \mathbf{Z}$ in time $O^*(2^{|U|} \cdot \log m)$ where m is a positive integer such that for all $S \subseteq U$, it holds that $|f(S)| \leq m$. The objective of this exercise is to show that the ranked version of the zeta transform can also be computed within this time. As usual, we assume that $f(S)$ for all $S \subseteq U = [n]$ can be computed in time $O^*(2^n \cdot \log m)$ using space $O^*(2^n \cdot \log m)$. The steps of the fast algorithm for the ranked zeta transform are as follows:

1. First compute the values of $f(S)$ for all $S \subseteq U$ and store them in a table of size $O^*(2^n \cdot \log m)$.
2. For $0 \leq i \leq n$, define $\zeta_k^{(i)}(f)$ as follows: for all $S \subseteq [n]$,

$$\zeta_k^{(i)}(f)(S) = \sum_{\substack{X \subseteq S; |X| = k \\ S \setminus [i] \subseteq X}} f(X).$$

   What do $\zeta_k^{(0)}(f)(S)$ and $\zeta_k^{(n)}(f)(S)$ represent?

3. If $i \in S$, show that
$$\zeta_k^{(i)}(f)(S) = \zeta_k^{(i-1)}(f)(S \setminus i) + \zeta_k^{(i-1)}(f)(S).$$

4. Now write down a recurrence for $\zeta_k^{(i)}(f)(S)$ in terms of $\zeta_k^{(i-1)}(f)(S\setminus i)$ and $\zeta_k^{(i-1)}(f)(S)$.

5. What is the time and space taken to compute $\zeta_k^{(i)}(f)(S)$ for all i, k, and S?

Homework Assignment H30 (10 Points) Counting k-Partitions. Let U be a set of n elements and let $\mathcal{F}$ be a family of subsets of U. A k-partition of the set system $(U, \mathcal{F})$ is a collection $S_1, \ldots, S_k$ of k sets from the family $\mathcal{F}$ such that $S_i \cap S_j = \emptyset$ for all $i \neq j$ and $\bigcup_{i=1}^{k} S_i = U$. Show that the number of k-partitions of a set system can be written as the iterated k-fold convolution of an appropriate function. This will show that an algorithm for computing the convolution of two functions may be used to count the number of k-partitions of a set system.

Homework Assignment H31 (10 Points) The Möbius and Zeta Transform. In class we saw that the Möbius transform of the zeta transform of a function $f\colon 2^U \to \mathbf{Z}$ is the function f itself. That is, for all $S \subseteq U$
$$\mu(\zeta(f))(S) = \sum_{X \subseteq S} (-1)^{|S\setminus X|} \zeta(f)(X) = f(S).$$
Show that the converse also holds, that is, the zeta transform of the Möbius transform of f is the function f itself.

Solution To prove that $\zeta(\mu(f)) = f$, it is sufficient to show that
$$f(S) = \sum_{X \subseteq S} \mu(f)(X) \tag{1}$$
for all $S \subseteq U$. By definition of the Möbius transform, the rhs of (1) may be written as
$$\begin{aligned} \sum_{X \subseteq S} \mu(f)(X) &= \sum_{X \subseteq S} \sum_{Y \subseteq X} (-1)^{|X\setminus Y|} f(Y) \\ &= \sum_{Y \subseteq S} \left(\sum_{\substack{X \subseteq S \\ Y \subseteq X}} (-1)^{|X\setminus Y|} \right) f(Y). \end{aligned}$$
The outer sum is over all subsets Y of S while the inner sum is over all supersets X of Y. Let $|Y| = p$ and $|S| = k$. If $k > p$ then the inner sum evaluates to
$$\sum_{i=0}^{k-p} (-1)^i \binom{k-p}{i} = (1-1)^{k-p} = 0.$$
If $p = k$ then the inner sum evaluates to 1 and the rhs of (1) is $f(S)$.

Homework Assignment H32 (10 Bonus Points) Fast Ranked Möbius Transform. Given a function $f\colon 2^{[n]} \to \mathbf{Z}$, show that the ranked Möbius transform of f can be computed in time $O^*(2^n \cdot \log m)$ time, where $m \in \mathbf{N}$ is such that for all $S \subseteq [n]$, it holds that $|f(S)| \leq m$. As usual, assume that for all $S \subseteq [n]$ one can compute $f(S)$ in time $O^*(2^n \cdot \log m)$ and space $O^*(2^n \cdot \log m)$.